



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GEOMETRY.

167. Proposed by JOHN M. QUINN, Professor of Mathematics, High School, Warren, Pa.

If at the vertex of an isosceles triangle one of whose basal vertices is pivoted and the other free to move in a straight line, a rhombus be pivoted with sides parallel to the sides of the triangle, the locus of every point on the rhombus except the one which is its intersection with the fixed side of the triangle is an ellipse.

Solution by the PROPOSER.

Notation. Let X and Y be rectangular axes; θ the angle BAX ; m equal segments of the sides of the triangle and the rhombus; $AB=a$, $BD=y$, and $AD=x$, and YBA an isosceles triangle with one basal vertex pivoted at A . The other basal vertex is free to move along the line AY .

To prove that the locus of any point as P is an ellipse.

PROOF. $y/a = \sin\theta$. $\therefore y^2/a^2 = \sin^2\theta$.


$x = a\cos\theta - 2m\cos\theta = \cos\theta(a - 2m)$.

$$\frac{x}{a-2m} = \cos\theta, \quad \frac{x^2}{(a-2m)^2} = \cos^2\theta.$$

$$\therefore \frac{y^2}{a^2} + \frac{x^2}{(a-2m)^2} = \sin^2\theta + \cos^2\theta = 1. \quad \therefore \text{the locus of } p \text{ is an ellipse.}$$

Similarly for any other point *mutatis mutandis*.

Q. E. D.



168. Proposed by MISS GUBELMAN, Student Southern Illinois State University, Carbondale, Ill.

To draw a perpendicular to one side of a triangle dividing it into two equivalent parts.

Solution by the PROPOSER.

1. Let ABC be the triangle. Draw the median AD and the perpendicular AE . Construct BX such that $BX^2 = BD \cdot BE$. Draw the perpendicular XY .

$$\therefore \triangle BXY : \triangle BAE = BX^2 : BE^2.$$


Similar triangles $= BD \times BE = BE^2 = BD : BE$.

Also $\triangle BAD : \triangle BAE = BD : BE$, having equal altitudes.

$$\therefore \triangle BXY : \triangle BAE = \triangle BAD : \triangle BAE.$$

$$\therefore \triangle BXY = \triangle BAD. \quad \text{But } \triangle BAD = \frac{1}{2} \triangle ABC. \quad \text{Median.}$$

$$\therefore \triangle BXY = \frac{1}{2} \triangle ABC. \quad \therefore XY \text{ is the required perpendicular.}$$



Also solved by G. B. M. ZERR, DANIEL B. NORTHRUP, L. C. WALKER, J. SCHEFFER, H. C. WHITAKER, H. B. PENHOLLOW, and P. W. WEBBER.

169. Proposed by S. F. NORRIS, Professor of Astronomy and Mathematics, Baltimore City College, Baltimore, Md.

Theorem. Two quadrilaterals having three sides of the one equal to the three corresponding sides of the other, each to each, and the two corresponding angles adjacent to the unknown sides equal, each to each, are equal figures. [From Olney's Geometry, Section VIII, Proposition XIV].

1. Required proof. 2. Is this proposition found in any other text-book of Geometry?